

ON WEAK*-CONVERGENCE IN THE HARDY SPACE H^1 OVER SPACES OF HOMOGENEOUS TYPE

HA DUY HUNG AND LUONG DANG KY *

ABSTRACT. Let $\mathcal{X}$ be a complete space of homogeneous type. In this note, we prove that the weak*-convergence is true in the Hardy space $H^1(\mathcal{X})$ of Coifman and Weiss.

1. INTRODUCTION

Let d be a quasi-metric on a set $\mathcal{X}$, that is, d is a nonnegative function on $\mathcal{X} \times \mathcal{X}$ satisfying

- $d(x, y) = d(y, x)$,
- $d(x, y) > 0$ if and only if $x \neq y$,
- there exists a constant $\kappa_1 \geq 1$ such that for all $x, y, z \in \mathcal{X}$,

$$(1.1) \quad d(x, z) \leq \kappa_1(d(x, y) + d(y, z)).$$

A trip $(\mathcal{X}, d, \mu)$ is called a *space of homogeneous type* in the sense of Coifman and Weiss if μ is a regular Borel measure satisfying *doubling property*, i.e. there exists a constant $\kappa_2 > 1$ such that for all $x \in \mathcal{X}$ and $r > 0$,

$$(1.2) \quad \mu(B(x, 2r)) \leq \kappa_2 \mu(B(x, r)).$$

In this paper, we always assume that $(\mathcal{X}, d, \mu)$ is a complete space of homogeneous type, $\mu(\mathcal{X}) = \infty$ and $0 < \mu(B) < \infty$ for any ball $B \subset \mathcal{X}$.

Recall (see [3]) that a function a is called an H^1 -atom related to the ball $B \subset \mathcal{X}$ if

- $\text{supp } a \subset B$;
- $\|a\|_{L^\infty(\mathcal{X})} \leq \mu(B)^{-1}$;
- $\int_{\mathcal{X}} a(x) d\mu(x) = 0$.

The Hardy space $H^1(\mathcal{X})$ is defined as the set of all $f = \sum_{j=1}^{\infty} \lambda_j a_j$ with $\{a_j\}_{j=1}^{\infty}$ are H^1 -atoms and $\{\lambda_j\}_{j=1}^{\infty} \subset \mathbb{C}$ is such that $\sum_{j=1}^{\infty} |\lambda_j| < \infty$. The norm on $H^1(\mathcal{X})$ is then defined by

$$\|f\|_{H^1(\mathcal{X})} := \inf \left\{ \sum_{j=1}^{\infty} |\lambda_j| : f = \sum_{j=1}^{\infty} \lambda_j a_j \right\}.$$

2010 *Mathematics Subject Classification.* 42B30.

Key words and phrases. H^1 , BMO, VMO, spaces of homogeneous type.

*Corresponding author.

It is well-known (see [3]) that the dual space of $H^1(\mathcal{X})$ is $BMO(\mathcal{X})$ the space of all locally integrable functions f with

$$\|f\|_{BMO(\mathcal{X})} := \sup_B \frac{1}{|B|} \int_B \left| f(x) - \frac{1}{|B|} \int_B f(y) d\mu(y) \right| d\mu(x) < \infty,$$

where the supremum is taken over all balls $B \subset \mathcal{X}$. Furthermore, $H^1(\mathcal{X})$ is itself the dual space of $VMO(\mathcal{X})$ the closure in $BMO(\mathcal{X})$ norm of the set $C_c(\mathcal{X})$ of all continuous functions with compact support.

The aim of the present paper is to establish the following.

Theorem 1.1. *Suppose that $\{f_n\}_{n=1}^\infty$ is a bounded sequence in $H^1(\mathcal{X})$, and that $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ for almost every $x \in \mathcal{X}$. Then, $f \in H^1(\mathcal{X})$ and $\{f_n\}_{n=1}^\infty$ weak*-converges to f , that is, for every $\varphi \in VMO(\mathcal{X})$, we have*

$$\lim_{n \rightarrow \infty} \int_{\mathcal{X}} f_n(x) \varphi(x) d\mu(x) = \int_{\mathcal{X}} f(x) \varphi(x) d\mu(x).$$

It should be pointed out that, when $\mathcal{X} = \mathbb{R}^d$, Theorem 1.1 was proved firstly by Jones and Journé [5] when trying to answer a question of Lions and Meyer [1]. When $\mathcal{X}$ is a norm space of homogeneous type, Theorem 1.1 was established by Grafakos and Rochberg [4].

2. PROOF OF THEOREM 1.1

Let M be the classical Hardy-Littlewood maximal function. The following result is well-known (see [3]).

Lemma 2.1. *There exists a constant $C_1 > 0$ such that*

$$\mu(\{x \in \mathcal{X} : Mg(x) > \lambda\}) \leq C_1 \frac{1}{\lambda} \int_{\mathcal{X}} |g(x)| d\mu(x)$$

for all $g \in L^1(\mathcal{X})$ and $\lambda > 0$.

By a standard argument (cf. [2, 4]), we get the following lemma.

Lemma 2.2. *There exists a constant $C_2 > 0$ such that, for any Borel set E ,*

$$\|\log(M(\chi_E))\|_{BMO(\mathcal{X})} \leq C_2.$$

Proof of Theorem 1.1. Without loss of generality, we can assume that $\|f_n\|_{H^1} \leq 1$ for all $n \geq 1$. Since the Fatou's lemma, we see that $\|f\|_{L^1(\mathcal{X})} \leq 1$. By this and a standard function theoretic argument, it suffices to show that

$$(2.1) \quad \lim_{n \rightarrow \infty} \int_{\mathcal{X}} f_n(x) \varphi(x) d\mu(x) = \int_{\mathcal{X}} f(x) \varphi(x) d\mu(x)$$

for all $\varphi \in C_c(\mathcal{X})$ with $\|\varphi\|_{L^1(\mathcal{X})}, \|\varphi\|_{L^\infty(\mathcal{X})} \leq 1$.

Fix $\varepsilon > 0$. As $f \in L^1(\mathcal{X})$, there exists a positive number β such that $\int_E |f| d\mu < \varepsilon$ for any Borel set A satisfying $\mu(A) < \beta$. By $\varphi \in C_c(\mathcal{X})$ and [6, Theorems 2 and 3], there exists $\alpha \in (0, \frac{\beta}{C_1 \varepsilon})$, where the constant C_1 is as in Lemma 2.1, such that if B is a ball in $\mathcal{X}$ satisfying $\mu(B) < \alpha$, then

$$(2.2) \quad |\varphi(x) - \varphi(y)| < \varepsilon$$

for all $x, y \in B$. On the other hand, by the Egorov's theorem and the regularity of μ , there exists an open set $E \subset \mathcal{X}$ such that $\mu(E) < \alpha \varepsilon e^{-\frac{1}{\varepsilon}}$ and $f_n \rightarrow f$ uniformly on $\text{supp } \varphi \setminus E$. Define $\tau := \max\{0, 1 + \varepsilon \log(M(\chi_E))\}$. It is clear that $0 \leq \tau \leq 1$ and $\tau(x) = 1$ for all $x \in E$. We now claim that

$$(2.3) \quad \|\varphi \tau\|_{BMO(\mathcal{X})} \leq (2 + 2C_1 + 3C_2)\varepsilon,$$

where the constant C_2 is as in Lemma 2.2.

Assume that (2.3) holds for a moment. Since $f_n \rightarrow f$ uniformly on $\text{supp } \varphi \setminus E$, there exists $n_0 \in \mathbb{N}$ such that $\|f_n - f\|_{L^\infty(\text{supp } \varphi \setminus E)} < \varepsilon$ for all $n \geq n_0$. Applying Lemma 2.1 with $g = \chi_E$ and $\lambda = e^{-\frac{1}{\varepsilon}}$, we obtain that

$$\mu(\text{supp } \tau) \leq C_1 \mu(E) e^{\frac{1}{\varepsilon}} < C_1 \alpha \varepsilon < \beta.$$

Therefore, for any $n \geq n_0$,

$$\begin{aligned} & \left| \int_{\mathcal{X}} (f_n - f) \varphi d\mu \right| \\ & \leq \left| \int_{\text{supp } \varphi \setminus E} (f_n - f) \varphi (1 - \tau) d\mu \right| + \left| \int_{\text{supp } \tau} f \varphi \tau d\mu \right| + \left| \int_{\mathcal{X}} f_n \varphi \tau d\mu \right| \\ & \leq \|f_n - f\|_{L^\infty(\text{supp } \varphi \setminus E)} \|\varphi\|_{L^1(\mathcal{X})} + \|\varphi\|_{L^\infty(\mathcal{X})} \int_{\text{supp } \tau} |f| d\mu + \|f_n\|_{H^1(\mathcal{X})} \|\varphi \tau\|_{BMO(\mathcal{X})} \\ & \leq \varepsilon + \varepsilon + (2 + 2C_1 + 3C_2)\varepsilon = (4 + 2C_1 + 3C_2)\varepsilon. \end{aligned}$$

This proves that (2.1) holds.

Let us now show (2.3). Let B be an arbitrary ball in $\mathcal{X}$. If $\mu(B) \geq \alpha$, then

$$\begin{aligned} \frac{1}{\mu(B)} \int_B \left| \varphi(x) \tau(x) - \frac{1}{\mu(B)} \int_B \varphi(y) \tau(y) d\mu(y) \right| d\mu(x) & \leq 2 \frac{1}{\mu(B)} \int_B |\varphi \tau| d\mu \\ & \leq 2 \frac{1}{\alpha} \|\varphi\|_{L^\infty(\mathcal{X})} \mu(\text{supp } \tau) \\ & \leq 2C_1 \varepsilon. \end{aligned}$$

Otherwise, by (2.2) and Lemma 2.2, we have

$$\begin{aligned}
& \frac{1}{\mu(B)} \int_B \left| \varphi(x)\tau(x) - \frac{1}{\mu(B)} \int_B \varphi(y)\tau(y)d\mu(y) \right| d\mu(x) \\
& \leq 2 \frac{1}{\mu(B)} \int_B \left| \varphi(x)\tau(x) - \frac{1}{\mu(B)} \int_B \varphi(y)d\mu(y) \frac{1}{\mu(B)} \int_B \tau(y)d\mu(y) \right| d\mu(x) \\
& \leq 2\|\tau\|_{L^\infty(\mathcal{X})} \sup_{x,y \in B} |\varphi(x) - \varphi(y)| + 2\|\varphi\|_{L^\infty(\mathcal{X})} \|\tau\|_{BMO(\mathcal{X})} \\
& \leq 2\varepsilon + 2\frac{3}{2}\|\varepsilon \log(M(\chi_E))\|_{BMO(\mathcal{X})} \leq (2 + 3C_2)\varepsilon,
\end{aligned}$$

which proves that (2.3) holds. $\square$

Acknowledgements. The paper was completed when the second author was visiting to Vietnam Institute for Advanced Study in Mathematics (VIASM). He would like to thank the VIASM for financial support and hospitality.

REFERENCES

- [1] R. R. Coifman, P.-L. Lions, Y. Meyer and S. Semmes, Compensated compactness and Hardy spaces. J. Math. Pures Appl. (9) 72 (1993), no. 3, 247–286.
- [2] R. R. Coifman and R. Rochberg, Another characterization of BMO. Proc. Amer. Math. Soc. 79 (1980), no. 2, 249–254.
- [3] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis. Bull. Amer. Math. Soc. 83 (1977), no. 4, 569–645.
- [4] L. Grafakos and R. Rochberg, Compensated compactness and the Heisenberg group. Math. Ann. 301 (1995), no. 3, 601–611.
- [5] P. W. Jones and J.-L. Journé, On weak convergence in $H^1(\mathbf{R}^d)$. Proc. Amer. Math. Soc. 120 (1994), no. 1, 137–138.
- [6] R. A. Macías and C. Segovia, Lipschitz functions on spaces of homogeneous type. Adv. in Math. 33 (1979), no. 3, 257–270.

Ha Duy Hung

High School for Gifted Students, Hanoi National University of Education, 136 Xuan Thuy, Hanoi, Vietnam

E-mail: hunghaduy@gmail.com

Luong Dang Ky

Department of Mathematics, University of Quy Nhon, 170 An Duong Vuong, Quy Nhon, Binh Dinh, Vietnam

E-mail: luongdangky@qnu.edu.vn